

Universal high-frequency transport in perfect photonic crystals

Chushun Tian^{1,2} and Luwei Zhou³

¹ *Institut für Theoretische Physik, Universität zu Köln, Köln, 50937, Germany*

² *Center for Nonlinear Studies, Hongkong Baptist University, Kowloon Tong, Hongkong*

³ *National key laboratory of surface physics and Department of physics, Fudan University, Shanghai, 200433, P. R. China*

(Dated: December 15, 2008)

The light scattering in the periodic dielectric cylinder array is studied. We analytically calculate the diffusive-ballistic transport crossover and find the weak localization superimposing on it. Possible experimental observations are analyzed.

PACS numbers: 42.25.Bs, 05.60.-k

Introduction—Experimental studies [1, 2] are offering new boost to interest in transport of optical systems. A traditional subject is the light scattering in fully [2] or partially [1, 3] disordered media. Irrespective of the absence/presence of the periodic background disorders turn out to be essential in both kinds of materials. There, the Anderson localization of light originates at the multiple disorder scattering. The latter leads to the enhanced constructive interference between two counterpropagating optical paths and, thus, suppresses the light diffusion.

For perfect periodic dielectric materials (namely photonic crystals) the low-frequency transport is protocolled by sample-specific band structures, the analysis of which is nowadays a well-established industry. By contrast, the high-frequency transport in perfect photonic crystals is a largely unexplored subject which might find practical applications. There, localization and periodicity may interplay with each other. Theoretically, no localization signatures have been found for periodic quantum mapping [4], which might suggest that localization is incompatible with perfect periodicity. Contrary to this, for realistic quantum particle motion in some perfect crystals, the weak localization correction to the classical diffusion coefficient is analytically shown [5]. Experimentally, to detect localization signals in periodic structures, if any, turns out to be extremely difficult. Indeed, in usual systems (e.g. electrons) ubiquitous particle interactions complicate analysis of wave interference effects such as (weak) localization. Although this problem is circumvented in optical systems, traditional transmission/reflection measurements do not allow directly probing the dynamical (bulk) diffusion coefficient.

The recent experiment [1] offers the very opportunity to study the diffusive-ballistic transport crossover. The experimental scope consists of two crucial steps: (i) to fabricate the dielectric material which is uniform in the longitudinal direction (but might be arbitrary in the transverse plane) and, (ii) to launch a probe laser beam, parallel to the longitudinal direction, into the material (Fig. 1). Then, the dynamical process of the dispersion in the transverse plane is recorded by the intensity profile at different propagating distances, which allows the di-




FIG. 1: A narrow input probe beam (green spot)—perpendicular to the plane—is launched into the dielectric cylinder array and expands. The “time” axis points inwards.

rect measurement of the dynamical diffusion coefficient. It is the purpose of this letter to analytically study the crossover in perfect photonic crystals and to show that weak localization superimposes on this crossover. Furthermore, we examine the possibility of confirming these predictions within this experimental scope.

Qualitative results based on band structure—To be specific we will consider the following photonic crystals. Dielectric cylinders of a radius r_0 , uniform in the “longitudinal” (z -) direction, are periodically embedded into the host material of a refractive index n_0 forming a triangular lattice in the “transverse” (x - y) plane with a lattice constant a (Fig. 1). For this lattice let us choose the coordinate system as $(x, y) = \mathbf{l} + \mathbf{r}$, where $\mathbf{l} \in \mathfrak{B}$ (the Bravais lattice) is chosen to be the cylinder center, and $\mathbf{r}$ specifies the position inside the unit cell. A probe beam of frequency Ω_0 , parallel to the z -axis, is launched into the host dielectric material, then expands as propagates along the z -direction. The transverse wave number $k_\perp$ is inversely proportional to the initial width of the beam, which is narrow enough such that $k_\perp^{-1} \ll \min(a - 2r_0, r_0)$.

Then, the propagation of the scalar field (for simplicity) denoted as $E(x, y, z)$ inside the photonic crystal is described by the Helmholtz equation (The light velocity in the vacuum is set to be unity.): $[\nabla^2 +$

$\Omega_0^2 n^2(x, y)]E(x, y, z) = 0$ with $n(x, y)$ the refractive index profile. Introduce the slow-varying envelope $\psi(x, y; z)$ such that $E(x, y, z) \equiv \psi(x, y; z) e^{i\Omega_0 n_0 z}$ and insert it into the Helmholtz equation. In the paraxial limit $|\partial_z^2 \psi| \ll \Omega_0 n_0 |\partial_z \psi|$ the second order derivative may be ignored [1], consequently we arrive at the Schrödinger equation:

$$\left\{ i \partial_z + \frac{1}{2\Omega_0 n_0} (\partial_x^2 + \partial_y^2) - V(x, y) \right\} \psi(x, y; z) = 0, \quad (1)$$

where $V(x, y) \equiv -\frac{\Omega_0}{2n_0} [n^2(x, y) - n_0^2]$. This shows that the beam propagates in the manner fully analogous to the two-dimensional quantum particle motion in the “potential” $V(x, y)$, and z plays the role of the usual time.

In periodic potentials exists the band structure $\epsilon_n(\mathbf{k})$, where n is the band index and $\mathbf{k}$ is the quasimomentum in the first Brillouin zone ($\mathbb{B}.Z.$). As to be detailed below (for a large class of potentials [6]) the ray optics inside a unit cell is ergodic characterized by the positive Lyapunov exponent λ . Following the well-known Bohigas-Giannoni-Schmit (BGS) conjecture, then, given $\mathbf{k}$ (away from symmetry points or lines) the high-frequency bands display universal level fluctuations [7].

If a single band is excited the motion of (Bloch) photons is ballistic. However, because the beam initially is narrow enough it excites a number of high-frequency bands centered at $\epsilon = k_\perp^2 / (2\Omega_0 n_0)$ with an amount of $\sim \lambda / \Delta \gg 1$, where Δ is the mean level spacing. At early times the rays are successively bent by dielectric cylinders which, equivalently, may be viewed as that Bloch photons experience considerable inelastic scattering. Consequently, the beam dispersion (expansion), defined as $\sigma^2(z) \equiv \iint dx dy (x^2 + y^2) |\psi(x, y; z)|^2$, is dominated by the interband transition factor: $e^{i\{\epsilon_n(\mathbf{k}) - \epsilon_{n'}(\mathbf{k})\}z}$. Then, the large number of excited bands allows us to average this factor with respect to the universal level correlator, which is $\sim \{\epsilon_n(\mathbf{k}) - \epsilon_{n'}(\mathbf{k})\}^{-2}$ [8, 9]. Consequently, the normal diffusion, i.e., $\sigma^2(z) \sim z$ is recovered for early times $\lambda^{-1} \lesssim z \ll \Delta^{-1}$. At later times $z \gtrsim \Delta^{-1}$ individual bands are resolved and the interband transition becomes negligible. The transport, in turn, becomes ballistic, i.e., $\sigma^2(z) \sim \Delta z^2$ [9].

For intermediate times $\lambda^{-1} \ll z \lesssim \Delta^{-1}$ photons move in the regime of size $\sim \sqrt{D_{cl}/\Delta}$, where the periodic structure is not resolved and the dielectric cylinders thereby resemble random scatterers. As usual while two optical paths diffusively propagate they find a significant probability to form a common loop and counter-propagate along it. Such constructive interference suppresses the normal diffusion coefficient D_{cl} and the weak localization correction results, which is order of $(\nu D_{cl})^{-1}$ with ν the photon density of states. Accordingly, the

linear dispersion is suppressed. That is,

$$\sigma^2(z) \sim \begin{cases} z, & z \gtrsim \lambda^{-1}, \\ z \left(1 - \frac{\ln \lambda z}{\nu D_{cl}} \right), & \lambda^{-1} \ll z \lesssim \Delta^{-1}, \\ \Delta z^2, & z \gtrsim \Delta^{-1}. \end{cases} \quad (2)$$

We now turn to the detailed analysis of wave dynamics described by Eq. (1) and derive the rigorous results. Below a finite system composed of $\mathcal{N}$ unit cells will be considered with the periodic boundary condition implemented. In the final results the limit: $\mathcal{N} \rightarrow +\infty$ is set.

Field-theoretic formalism—First we present the general formalism used throughout this work. Consider the propagation on the lattice from $\mathbf{l}$ to $\mathbf{l}'$, described by the so-called two-point correlator: $\mathcal{Y}(\mathbf{l} - \mathbf{l}'; \omega) \equiv \mathcal{N} \iint d\mathbf{r} d\mathbf{r}' G_{\epsilon + \frac{\omega}{2}}^R(\mathbf{r} + \mathbf{l}, \mathbf{r}' + \mathbf{l}') G_{\epsilon - \frac{\omega}{2}}^A(\mathbf{r}' + \mathbf{l}', \mathbf{r} + \mathbf{l})$, noticing that $\mathbf{r}, \mathbf{r}'$ are the coordinates in the unit cell. Here $G^{R/A}$ is the retarded/advanced Green function of Eq. (1) in the frequency domain. The translational symmetry implies that $\mathcal{Y}$ depends merely on $\mathbf{l} - \mathbf{l}'$.

Following Ref. [10] the Green functions are written down explicitly in the Bloch basis $|n\mathbf{k}\rangle$. In doing so the wave dynamics is reduced into the one in a two-torus. For the latter we invoke the Wigner transform and pass to the phase space $\mathbb{T}$. The coordinate is denoted as $X \equiv (\mathbf{r}, \mathbf{p})$ (after appropriate rescaling) and satisfies $\int_{\mathbb{T}} dX = 1$. Then, it is a canonical procedure [8, 10, 11] to express the spatial Fourier transform of $\mathcal{Y}(\mathbf{l} - \mathbf{l}'; \omega)$, denoted as $\mathcal{Y}(\mathbf{q}, \omega)$, in terms of the functional integral over the supermatrix field $Q = T\Lambda T^{-1}$, where both Q and T depend on X . Moreover, $Q(X) = K C^T Q^T(\bar{X}) C K$ and $T^\dagger(X) = C T^T(\bar{X}) C^T$ with $\bar{X} \equiv (\mathbf{r}, -\mathbf{p})$. After tedious calculations we find (Throughout this work we are interested in the low-lying modes, i.e., $\mathbf{q} \rightarrow 0$.)

$$\mathcal{Y}(\mathbf{q}, \omega) = \frac{(\pi\nu)^2}{64\mathcal{N}} \sum_{\mathbf{k}_\pm} \iint_{\mathbb{T}} dX dX' e^{i\mathbf{q} \cdot (\mathbf{r} - \mathbf{r}')} \int DQ e^{-F[Q, \mathbf{a}]} \times \text{str} [k(1 + \Lambda)(1 - \tau_3)Q(X)k(1 - \Lambda)(1 - \tau_3)Q(X')] , \quad (3)$$

where $\tilde{\Sigma}_{\mathbf{k}_\pm} \equiv \Sigma_{\mathbf{k}_\pm \in \mathbb{B}.Z.} \delta_{\mathbf{k}_+ - \mathbf{k}_-, \mathbf{q}}$, str is the supertrace, and the action is

$$F[Q, \mathbf{a}] = \frac{\pi\nu}{2} \int_{\mathbb{T}} dX \text{str} \left\{ T\Lambda \hat{\mathcal{L}}_a T^{-1} + \frac{i\omega^+}{2} \Lambda Q \right\}. \quad (4)$$

Here all the constant supermatrices: Λ, τ_3, k, K and C , as well as the supermatrix fields Q and T are 8×8 matrices defined on the retarded/advanced, bosonic/fermionic and time-reversal sector (for the details of which we refer to Ref. [8]). In Eq. (3) $\hat{\mathcal{L}}_a = \mathbf{p} \cdot (\partial_{\mathbf{r}} + [i\mathbf{a}\tau_3, \cdot]) - \partial_{\mathbf{r}} V \cdot \partial_{\mathbf{p}}$ is the covariant Liouvillian, where the supermatrix $\mathbf{a} = \text{diag}(\mathbf{k}_+, \mathbf{k}_-)^{\text{ra}}$ with the superscript standing for the retarded/advanced sector. Eqs. (3) and (4) are the exact starting point of the succeeding analysis.

We remark that the coarse graining $\mathbb{T}$, with the element commensurating with the uncertainty intrinsic to

wave dynamics [12], suffices to regularize the field theory. (The results below, however, are insensitive to the details which, therefore, are not presented.) Indeed, the Liouvillian $\hat{\mathcal{L}}_0$ induces a deterministic flow on $\mathbb{T}$: $X \mapsto X(t) \equiv (\mathbf{r}(t), \mathbf{p}(t))$. The flow (i) has positive λ and is ergodic and, (ii) enjoys exponential decay of correlations and the central limiting theorem for sufficiently regular (Hölder-continuous) functions. They lead to singular hydrodynamic (steady) states [13] (and justify the application of the BGS conjecture in the qualitative discussions), as we also show below. Then, associated with the coarse graining $\mathbb{T}$ the Gibb's entropy production results which can be shown to be positive-definite [13], signaling a stable field theory.

Notice that properties (i) and (ii) follow directly from the choice of the potential V [6]. Indeed, the former is given by the Donnay-Liverani theorem [14]. To justify the latter we introduce the Birkhoff coordinate: (θ, φ) , where θ is the position of the incident ray at the dielectric cylinder boundary ($r = r_0$), and $\varphi \in [0, \pi]$ is the angle made between the incident ray and the tangential direction of the cylinder boundary. Then, we observe that the rotation function $\Delta\theta(\varphi)$ defined in [15], as well as its first-order derivative, is discontinuous at $\varphi = \frac{\pi}{2}$ while continuous at each interval. From the Bálint-Tóth theorem [14] property (ii) immediately follows.

Hydrodynamic relaxation and Gaspard nonequilibrium steady state—Let us start from the ray optics limit: $\Delta/\lambda \rightarrow 0$ and present a field-theoretic proof of the *deterministic* diffusion. The emergence of hydrodynamic relaxation from the Liouvillian dynamics is a long standing problem [16]. It has been exactly solved for the periodic Lorentz gases [13] as discussed here. To calculate Eq. (3) we employ the so-called rational parametrization: $T = 1 + iW$. Here $W\Lambda + \Lambda W = 0$ and, moreover, $W(X) = -KCW^T(\bar{X})C^TK$. Inserting it into the action F the perturbative expansion in W shows that W^2 is order of Δ/λ . Because of $\Delta/\lambda \rightarrow 0$ in the expansion of both F and the prefactor only the quadratic terms are to be kept. Upon fixing the $U(1)$ gauge the succeeding calculations are formally parallel to disordered systems [8], but, there is a crucial difference namely the resolvent: $\{-i\omega^+ + \hat{\mathcal{L}}_0 + \mathbf{p} \cdot i\mathbf{g}\}^{-1}$. Recall that, in disordered systems, this field propagator is replaced by the heat kernel recovering the diffusion. Upon passing to the time domain the resolvent, denoted as $\mathcal{D}_{\mathbf{g},t}$, solves

$$\mathcal{D}_{\mathbf{g},t} = e^{-t\hat{\mathcal{L}}_0} - \int_0^t d\tau e^{-(t-\tau)\hat{\mathcal{L}}_0} \mathbf{p} \cdot i\mathbf{g} \mathcal{D}_{\mathbf{g},\tau}. \quad (5)$$

For low-lying modes the functional equation (5) gives

$$\begin{aligned} \mathcal{D}_{\mathbf{g},t} = & \left\{ 1 - \int_0^t d\tau \mathbf{p}(-\tau) \cdot i\mathbf{g} \right. \\ & \left. - \int_0^t d\tau \int_0^{t-\tau} d\tau' \mathbf{g}\mathbf{g} : \mathbf{p}(-\tau)\mathbf{p}(-\tau-\tau') \right\} e^{-t\hat{\mathcal{L}}_0} \end{aligned} \quad (6)$$

with the second-order iteration. Combined with property (i) Eq. (6) shows that, given a generic initial distribution, for vanishing $\mathbf{g}$ it converges to the invariant microcanonical measure in the limit $t \rightarrow +\infty$. A small concentration gradient $i\mathbf{g}$ drives the system out of equilibrium:

$$\mathcal{D}_{\mathbf{g},t \rightarrow +\infty} = 1 - \int_0^{+\infty} d\tau \mathbf{p}(-\tau) \cdot i\mathbf{g} + o(\mathbf{g}^2), \quad (7)$$

where the second term is the singular microscopic nonequilibrium steady state found by Gaspard [13]. Furthermore, by inserting Eq. (6) into Eq. (3) and returning back to the frequency domain, we find $\mathcal{Y}(\mathbf{q}, \omega) = 2\pi\nu/(-i\omega + D_{cl}\mathbf{q}^2)$ and, with the help of property (ii), rigorously justify the Green-Kubo formula:

$$D_{cl} = \frac{1}{2} \int_{\mathbb{T}} dX \int_0^{+\infty} dt \mathbf{p}(t) \cdot \mathbf{p}(0). \quad (8)$$

Eqs. (7) and (8) confirm that the normal diffusion appears as the Pollicott-Ruelle resonance as shown earlier [13]. There, a completely different technique—the ζ -function theory—is used informing its far-reaching relation to the field-theoretic approach. In obtaining the diffusive two-point correlator the integration over X and X' is essential by which the scaling regime is recognized.

Weak localization—For $\Delta/\lambda \ll 1$ wave interference accounts for fluctuations. Keeping the W -expansion up to the fourth order the resolvent is modified to be $\{-i\omega^+ + \hat{\mathcal{L}}_0 + \delta\hat{\mathcal{L}} + \mathbf{p} \cdot i\mathbf{g}\}^{-1}$. Here the perturbed operator $\delta\hat{\mathcal{L}}$ is explicitly expressed as

$$\begin{aligned} \delta\hat{\mathcal{L}}_{XX'} = & -\frac{1}{\mathcal{N}} \sum_{\mathbf{g} \in \mathbb{B.Z.}} \int_0^{+\infty} dt e^{-i\omega^+ t} \langle \bar{X} | \mathcal{D}_{\mathbf{g},t} | X \rangle \\ & \times \{ \langle X | X' \rangle - \delta(X - X') \} \end{aligned} \quad (9)$$

which, in the ray optics limit, is nullified by the fine-grained phase space structure because the bases $\langle X |$ and $| X' \rangle$ satisfy $\langle X | X' \rangle = \delta(X - X')$. In wave dynamics such point-like structure is smeared by the uncertainty. Indeed, a famous theorem of periodic Lorentz gases suggests that the bases $| X \rangle / \langle X |$ must instead be constructed as a distribution supported by a small neighborhood of X and normalized to unity, with $\langle X | X' \rangle$ defined as the overlap between two coarse grained representing points X and X' [17]. (Notice that the point-like distribution cannot be spanned by these bases.) Furthermore, it was rigorously shown [17] that $| X \rangle$ relaxes to the microcanonical measure at some short time scale ($\gtrsim \lambda^{-1}$) when the support of $| X \rangle$ uniformly covers $\mathbb{T}$. Upon combined with Eq. (6) the matrix element $\langle \bar{X} | \mathcal{D}_{\mathbf{g},t} | X \rangle$, in turn, acquires a universal value while vanishes at shorter times $t \lesssim \lambda^{-1}$. (For chaotic cavities this fact was noticed by Müller and Altland [12].) Importantly, such an interference correction conserves the probability (obeying the so-called Ward identity) guaranteed by $\int_{\mathbb{T}} dX \{ \langle X | X' \rangle - \delta(X - X') \} = \int_{\mathbb{T}} dX' \{ \langle X | X' \rangle - \delta(X - X') \} = 0$.

Repeat the same procedure of Eqs. (5)-(8) taking into account the third order iteration. After tedious calculations we arrive at a similar diffusive two-point correlator but, crucially, D_{cl} is renormalized into

$$D_0(\omega) = D_{cl} \left[1 - \frac{1}{\pi\nu} \int \frac{d\mathbf{q}}{(2\pi)^2} \frac{1}{-i\omega + D_{cl}\mathbf{q}^2} \right] \quad (10)$$

for $\Delta \lesssim \omega \ll \lambda$. The second term fully agrees with earlier theoretical prediction for periodic Lorentz gases [5], and resembles the well-known weak localization correction of disordered systems. In the latter systems strong localization develops for lower ω . Nevertheless here such a scenario breaks down which we now come to study.

Diffusive-ballistic transport crossover—For $\omega \ll \lambda$ the zero mode (uniform component) of the T field, denoted as T_0 is established recognizing the microcanonical measure over $\mathbb{T}$. Therefore, we factorize T as $T = T_0 T_{>}$ and insert it into the action, where $T_{>}$ are the fluctuating component. With $T_{>}$ integrated out Eq. (3) gives

$$\mathcal{Y}(\mathbf{q} \rightarrow 0, \omega) = \frac{(\pi\nu)^2}{64N} \sum_{\mathbf{k}_{\pm}} \int DQ_0 e^{-\tilde{F}[Q_0, \mathbf{a}]} \times \text{str} [k(1 + \Lambda)(1 - \tau_3)Q_0 k(1 - \Lambda)(1 - \tau_3)Q_0], \quad (11)$$

where $Q_0 = T_0 \Lambda T_0^{-1}$ and the zero mode action: $\tilde{F}[Q_0, \mathbf{a}] = \frac{\pi\nu}{8} \text{str} \{ D^* [i\mathbf{a}\tau_3, Q_0]^2 + \frac{i\omega^+}{2} \Lambda Q_0 \}$. Here $D^* = \text{Re } D_0(\max\{\omega, \Delta\})$ with $\max\{\omega, \Delta\}$ playing the role of the infrared cutoff.

To proceed further we invoke the Efetov parametrization for Q_0 [8]. Then, the hydrodynamic (i.e., $\mathbf{a}$ -) expansion for Eq. (11) is performed and kept up to the second order. The remaining integral can be exactly done giving

$$\mathcal{Y}(\mathbf{q} \rightarrow 0, \omega) = \frac{2\pi\nu}{-i\omega + D(\omega)\mathbf{q}^2}, \quad \omega \ll \lambda, \quad D(\omega) = D^* \left[1 - \frac{\Delta^2}{2\pi\omega^2} (1 - e^{2i\pi\omega/\Delta}) \right]. \quad (12)$$

Strikingly, the diffusive-ballistic transport crossover is universal despite that the present unit cell drastically differs from a disordered one [18]. The fundamental structure difference at the unit cell level profoundly affects the ray optics and thereby the field theory construction. Notice that D_{cl} is now renormalized into D^* which is not reported for periodic disordered systems [18]. These low-lying modes are diffusive for sufficiently large frequencies, i.e., $\Delta \ll \omega \lesssim \lambda$, while ballistic in the opposite limit, i.e., $\omega \ll \Delta$ because of $D(\omega) \approx D_0(\Delta)\Delta/(-i\pi\omega)$.

Possible experimental observations—Eqs. (7), (8), (10) and (12) are the main results of this work. To confirm the universal diffusive-ballistic crossover might be within the reach of the experimental scope of Ref. [1] utilizing the optical induction technique [19]. The incident probe beam is of $10\mu\text{m}$ full-width at half-maximum. The refractive index is required to vary over a scale of $100\mu\text{m}$

which, we expect, might be possible by tuning the optical interference pattern. The output light intensity profile $I(x, y; z)$ then determines the dispersion through $\sigma^2(z) = \iint dx dy (x^2 + y^2) I(x, y; z)$. The latter is directly transferred into $D(\omega)$ by

$$\sigma^2(z) - \sigma^2(0) = \int \frac{d\omega}{\pi} \frac{1 - e^{-i\omega z}}{\omega^2} D(\omega), \quad (13)$$

giving Eq. (2) with Eq. (12) inserted. Finally, we stress that the universality of high-frequency transport relies on the normal diffusion in the ray optical limit. Thus, the predictions are applicable for a large class of periodic dielectric structures. In particular, the weak localization is stronger for the quasi-one-dimensional transverse plane.

We are grateful to M. Garst for a valuable contribution. We have also enjoyed discussions with A. Altland, J. R. Dorfman, P. Gaspard, F. Haake, and Z. Q. Zhang, and were encouraged by Bambi Hu. This work is supported by Transregio SFB 12 of the Deutsche Forschungsgemeinschaft, NNSF of China (No. 10334020 and 10574027), and partly by the Hongkong Baptist University.

-
- [1] T. Schwartz, *et. al.*, Nature **446**, 52 (2007).
 - [2] M. Stözer, *et. al.*, Phys. Rev. Lett. **96**, 063904 (2006).
 - [3] A. F. Koenderink, *et. al.*, Phys. Lett. A **268**, 104 (2000).
 - [4] D. K. Wójcik and J. R. Dorfman, Phys. Rev. Lett. **90**, 230602 (2003); Physica D **187**, 223 (2004).
 - [5] C. Tian and A. Larkin, Phys. Rev. Lett. **95**, 246601 (2005).
 - [6] The refractive index inside the cylinder is central symmetric and is denoted as $n(r)$, $r \in (0, r_0)$. It satisfies a) $V \in C^3((0, r_0))$, b) $h'(r) > 0$ for all $r \in (0, r_0)$, c) $\lim_{r \rightarrow 0} r^2 V(r) = 0$, $V(r_0) = 0$, $V'(r_0^-) < 0$, d) $\Omega(r) < 0$ and e) $a - 2r_0 > -2/V'(r_0^-)$. Here $h(r) \equiv r^2[k_{\perp}^2 - 2\Omega_0 n_0 V(r)]$ and $\Omega(r) \equiv rh'(r)/h(r)$. These conditions, together with the cylinder geometry are merely for technical reasons. Furthermore, a satisfies $2r_0 < a \leq \frac{4}{\sqrt{3}}r_0$ so that the mean free path is bounded from above.
 - [7] E. R. Mucciolo, *et. al.*, Phys. Rev. B **50**, 8245 (1994).
 - [8] K. B. Efetov, *Supersymmetry in disorder and chaos* (Cambridge, UK, 1997).
 - [9] C. Tian and M. Garst, unpublished.
 - [10] C. Tian, Pis'ma Zh. Eksp. Teor. Fiz. **86**, 651 (2007) [JETP Lett. **86**, 566 (2008)]; Phys. Rev. B **77**, 064205 (2008).
 - [11] A. V. Andreev, *et. al.*, Nucl. Phys. B **482**, 536 (1996).
 - [12] C. Tian, A. Kamenev, and A. Larkin, Phys. Rev. Lett. **93**, 124101 (2004); J. Müller and A. Altland, J. Phys. A **38**, 3097 (2005); J. Müller T. Micklitz, and A. Altland, Phys. Rev. E **76**, 056204 (2007).
 - [13] P. Gaspard, Phys. Rev. E **53**, 4379 (1996); J. R. Dorfman, P. Gaspard, and T. Gilbert, *ibid.*, **66**, 026110 (2002).
 - [14] V. Donnay and C. Liverani, Comm. Math. Phys. **135**, 267 (1991); P. Bálint and I. P. Tóth, *ibid.*, **243**, 55 (2003).
 - [15] The rotation function is defined as $\Delta\theta(\varphi) \equiv 2r_0 \cos \varphi \int_{\hat{r}(\varphi)}^{r_0} r^{-1} \{h(r) - (r_0 \cos \varphi)^2\}^{-1/2} dr$, where $\hat{r}(\varphi)$

is the minimal distance as the trajectory approaches the cylinder center, and $h(r)$ is defined in [6].

- [16] I. Prigogine, *Nonequilibrium statistical mechanics* (Wiley, New York, 1962).
- [17] L. A. Bunimovich and Ya. G. Sinai, Comm. Math. Phys. **78**, 247 (1980). The accurate numerical prefactor of λ^{-1}
- is difficult to compute and logarithmically depends on the size of the support of $|X\rangle$.
- [18] N. Taniguchi and B. L. Altshuler, Phys. Rev. Lett. **71**, 4031 (1993).
- [19] N. K. Efemidis, *et. al.*, Phys. Rev. E **66**, 046602 (2002).